

Find out how to access preview-only contentLook inside Get Access
Advances in Meteorology, Climatology and Atmospheric Physics

Springer Atmospheric Sciences 2013, pp 317-323

The Effects of Naturally Produced Dust Particles on Radiative Transfer

Abstract

Mineral dust has a profound effect on the radiative budget and energy distribution of the atmosphere. By absorbing and scattering the solar radiation aerosols reduce the amount of energy reaching the surface. In addition aerosols enhance the greenhouse effect by absorbing and emitting longwave radiation. Desert dust forcing exhibits large regional and temporal variability due to its short lifetime and diverse optical properties further complicate the quantification of the Direct Radiative Effect (DRE). The complexity of the above processes, indicate the need of an integrated approach in order to examine these impacts. To this end the radiative transfer module RRTMG has been incorporated into the framework of the SKIRON model. The updated system was used to perform a 6-year long simulation over the Mediterranean region. As it was found, the most profound effect dust clouds have in areas away from the sources is the surface cooling through the “shading” effect. The long wave radiation forcing below and above the dust cloud is considerable and drives changes in the tropospheric temperature. In general dust particles cause warming near the ground and at mid-tropospheric layers and at the same time cooling of the lower troposphere.

Page %P

Page 1

The Effects of Naturally Produced Dust Particles on Radiative Transfer

C. Spyrou, G. Kallos, C. Mitsakou, P. Athanasiadis, and C. Kalogeri

Abstract Mineral dust has a profound effect on the radiative budget and energy distribution of the atmosphere. By absorbing and scattering the solar radiation aerosols reduce the amount of energy reaching the surface. In addition aerosols enhance the greenhouse effect by absorbing and emitting longwave radiation. Desert dust forcing exhibits large regional and temporal variability due to its short lifetime and diverse optical properties further complicate the quantification of the Direct Radiative Effect (DRE). The complexity of the above processes, indicate the need of an integrated approach in order to examine these impacts. To this end the radiative transfer module RRTMG has been incorporated into the framework of the SKIRON model. The updated system was used to perform a 6-year long simulation over the Mediterranean region. As it was found, the most profound effect dust clouds have in areas away from the sources is the surface cooling through the “shading” effect. The long wave radiation forcing below and above the dust cloud is considerable and drives changes in the tropospheric temperature. In general dust particles cause warming near the ground and at mid-tropospheric layers and at the same time cooling of the lower troposphere.

1 Introduction

Large amounts of mineral dust are produced from arid and semi-arid areas under favorable conditions. Aerosols interact strongly with solar and terrestrial radiation in several ways (Tegen et al. 1996; Haywood et al. 2003; Yoshioka et al. 2005; IPCC 2007; Kallos et al. 2009), also known as “Direct Aerosol Effect – DRE” (IPCC 2007). By absorbing and scattering the solar radiation, aerosols reduce

C. Spyrou (✉) • G. Kallos • C. Mitsakou • P. Athanasiadis • C. Kalogeri
Atmospheric Modeling and Weather Forecasting Group, Department of Physics,
University of Athens, University Campus, Building Physics V, Athens, Greece
e-mail: cspir@mg.uoa.gr

No Body Text -- translate me!

Page 2

the amount of energy reaching the surface (Kaufman et al. 2005; Tegen 2003; Spyrou et al. 2010). Moreover, aerosols enhance the greenhouse effect by absorbing and emitting outgoing longwave radiation (Dufrense et al. 2001; Tegen 2003).

Aerosols act as a cloud condensation nuclei or ice nuclei, thus modifying the microphysical, microchemical, and, hence, optical properties of clouds (Levin et al. 2005; Solomos et al. 2011). Collectively changes in cloud processes due to aerosols are referred to as aerosol indirect effects. Finally, absorption of solar radiation by particles contributes to the reduction in cloudiness, a phenomenon referred to as the semi-direct effect. This occurs because absorbing aerosol warms the atmosphere, which changes the atmospheric stability, and reduces surface flux (IPCC 2007).

The magnitude of the feedback on the radiative transfer depends strongly on the optical properties of particles (single scattering albedo, asymmetry parameter, extinction efficiency), which in turn depend on the size, shape and refractive indexes of dust particles (Tegen 2003; Helmert et al. 2007). The mineral composition of the dust source areas (Tegen 2003), as well as the chemical composition and transformation of aerosols during their transportation (Astitha et al. 2010) are all factors on the optical intensity of dust. Furthermore the vertical distribution of dust, the presence of clouds and the albedo of the surface all contribute to the DRE (Sokolik and Toon 1996; Tegen and Lacis 1996; Liao and Seinfeld 1998; Helmert et al. 2007).


In the present study, the main objective is to quantify the contribution of dust particles on the DRE. To this end, the radiative transfer module – RRTMG (Mlawer et al. 1997; Oreopoulos and Barker 1999; Iacono et al. 2003; Pincus et al. 2003; Barker et al. 2003; Clough et al. 2005; Morcrette et al. 2008; Iacono et al. 2008) has been incorporated into the framework of the SKIRON/DUST model. The SKIRON/Dust system has already the capability to adequately simulate the desert dust cycle using state of the art parameterizations for the uplift, transportation and deposition of aerosols (Spyrou et al. 2010). The addition of the RRTMG radiative scheme has made it possible to model and study the effects of desert dust particles to the radiation balance of the atmosphere.

2 Experimental Design

In order to study the effects of desert dust particles to the radiation balance of the atmosphere, simulations for a 6-year period (2002–2007) were carried out with the implementation of the improved SKIRON/Dust modeling system.

The model was integrated over an extended area that covers the European continent, the Mediterranean Sea and northern Africa, as well as a major part of Middle East and Turkey. The horizontal grid increment was 0.24° and in the vertical direction 38 levels were used stretching from the ground surface up to 20 km. A high resolution reanalysis dataset (horizontal resolution of 0.15°) that has been developed in the framework of CIRCE project (Climate Change and Impact

No Body Text -- translate me!



Citations

Within this Chapter

1. Introduction
2. Experimental Design
3. Results
4. Conclusions
5. References
6. References

Related Content



References (24)

1. Astitha M, Kallos G, Spyrou C, O'Hirok W, Lelieveld J, Denier van der Gon HAC (2010) Modelling the chemically aged and mixed aerosols over the eastern central Atlantic Ocean – potential impacts. *Atmos Chem Phys* 10:5797–5822. doi:10.5194/acp-10-5797-2010 CrossRef
2. Barker HW, Pincus R, Morcrette JJ, The Monte-Carlo Independent Column Approximation (2003) Application within large-scale models. In: Proceedings of the GCSS/ARM workshop on the representation of cloud systems in large-scale models, Kananaskis, Alberta, Canada, 10pp
3. Clough SA, Shephard MW, Mlawer EJ, Delamere JS, Iacono MJ, Cady-Pereira K, Boukabara S, Brown PD (2005) Atmospheric radiative transfer modeling: a summary of the AER codes. *J Quant Spectrosc Radiat Transf* 91:233–244 CrossRef
4. Dufrense JL, Gautier C, Ricchiazzi P (2001) Longwave scattering of mineral aerosols. *J Atmos Sci* 59:959–1966
5. Haywood JM et al (2003) Radiative properties and direct Radiative effect of Saharan dust measured by the C-130 aircraft during SHADE: 1. Solar spectrum. *J Geophys Res* 108(D18):8577. doi:10.1029/2002JD002687 CrossRef
6. Helmert J, Heinold B, Tegen I, Hellmuth O, Wendisch M (2007) On the direct and semidirect effects of Saharan dust over Europe: a modelling study. *J Geophys Res* 112. doi:10.1029/2006JD007444
7. Iacono MJ, Delamere JS, Mlawer EJ, Clough SA (2003) Evaluation of upper tropospheric water vapor in the NCAR community climate model (CCM3) using modeled and observed HIRS radiances. *J Geophys Res* 108(D2):4037. doi:10.1029/2002JD002539 CrossRef
8. Iacono MJ, Delamere JS, Mlawer EJ, Shephard MW, Clough SA, Collins WD (2008) Radiative forcing by long-lived greenhouse gases: calculations with the AER radiative transfer models. *J Geophys Res* 113:D13103. doi:10.1029/2008JD009944 CrossRef
9. Intergovernmental Panel on Climate Change (IPCC) (2007), *Climate change 2007: the physical science basis*, Cambridge University Press, UK
10. Kallos G, Spyrou C, Astitha M, Mitsakou C, Solomos S, Kushta J, Pytharoulis I, Katsafados P, Mavromatidis E, Papantoniou N, Vlastou G (2009) Ten-year operational dust forecasting – recent model development and future plans. *IOP Conf Ser Earth Environ Sci* 7(2009). doi:10.1088/1755-1307/7/1/012012
11. Kaufman YJ, Koren I, Remer LA, Rosenfeld D, Rudich Y (2005) The effect of smoke, dust

- and pollution aerosol on shallow cloud development over the Atlantic 160 Ocean. *Proc Natl Acad Sci USA* 102:11207–11212 CrossRef
12. Levin Z, Teller A, Ganor E, Yin Y (2005) On the interactions of mineral dust, sea-salt particles and clouds: a measurement and modelling study from the Mediterranean Israeli Dust Experiment campaign. *J Geophys Res* 110:D20202. doi:10.1029/2005JD005810 CrossRef
 13. Liao H, Seinfeld JH (1998) Radiative forcing by mineral dust aerosols: sensitivity to key variables. *J Geophys Res* 103(D):31637–31645 CrossRef
 14. Mlawer EJ, Taubman SJ, Brown PD, Iacono MJ, Clough SA (1997) RRTM, a validated correlated-k model for the longwave. *J Geophys Res* 102:16,663–16,682 CrossRef
 15. Morcrette JJ, Barker HW, Cole JNS, Iacono MJ, Pincus R (2008) Impact of a new radiation package, McRad, in the ECMWF integrated forecast system. *Mon Weather Rev* 136(12):4773–4798, doi: 10.1175/2008MWR2363.1
 16. Oreopoulos L, Barker HW (1999) Accounting for subgrid-scale cloud variability in a multi-layer 1-D solar radiative transfer algorithm. *Q J R Meteor Soc* 125:301–330
 17. Pincus R, Barker HW, Morcrette JJ (2003) A fast, flexible, approximate technique for computing radiative transfer in inhomogeneous clouds. *J Geophys Res* 108(D13):4376. doi:10.1029/2002JD003322 CrossRef
 18. Sokolik IN, Toon OB (1996) Direct radiative forcing by anthropogenic airborne mineral aerosols. *Nature* 381:681–683 CrossRef
 19. Solomos S, Kallos G, Kushta J, Astitha M, Tremback C, Nenes A, Levin Z (2011) An integrated modeling study on the effects of mineral dust and sea salt particles on clouds and precipitation. *Atmos Chem Phys* 11:873–892. doi:10.5194/acp-11-873-2011 CrossRef
 20. Spyrou C, Mitsakou C, Kallos G, Louka P, Vlastou G (2010) An improved limited area model for describing the dust cycle in the atmosphere. *J Geophys Res* 115:D17211. doi:10.1029/2009JD013682 CrossRef
 21. Tegen I (2003) Modeling the mineral dust aerosol cycle in the climate system. *Q Sci Rev* 22:1821–1834 CrossRef
 22. Tegen I, Lacis AA (1996) Modeling of particle size distribution and its influence on the radiative properties of mineral dust aerosol. *J Geophys Res* 101:19,237–19,244 CrossRef
 23. Wilks DS (1995) *Statistical methods in the atmospheric sciences*, Academic Press NY, pp 233–277
 24. Yoshioka M, Mahowald N, Dufresne JL, Luo C (2005) Simulation of absorbing aerosol indices for African dust. *J Geophys Res* 110:D18S17. doi:10.1029/2004JD005276 CrossRef

Title

The Effects of Naturally Produced Dust Particles on Radiative Transfer

Book Title

Advances in Meteorology, Climatology and Atmospheric Physics

Book Part

Part I

Pages

pp 317-323

Copyright

2013

DOI

10.1007/978-3-642-29172-2_45

Print ISBN

978-3-642-29171-5

Online ISBN

978-3-642-29172-2

Series Title

Springer Atmospheric Sciences

Series ISSN

2194-5217

Publisher

Springer Berlin Heidelberg

Copyright Holder

Springer-Verlag Berlin Heidelberg

Additional Links

- [About this Book](#)



Topics

- [Meteorology/Climatology](#)
- [Natural Hazards](#)
- [Environmental Monitoring/Analysis](#)
- [Atmospheric Protection/Air Quality Control/Air Pollution](#)
- [Environmental Health](#)
- [Climate Change](#)

eBook Packages

- [eBook Package english full Collection](#)
- [eBook Package english Earth & Environmental Science](#)

Editors

- [Costas G. Helmis](#)  (ID1)
- [Panagiotis T. Nastos](#)  (ID2)

Editor Affiliations

- ID1. , Department of Physics, University of Athens
- ID2. , Geography and Climatology, University of Athens

Authors

- C. Spyrou ⁽¹⁾
- G. Kallos ⁽¹⁾
- C. Mitsakou ⁽¹⁾
- P. Athanasiadis ⁽¹⁾
- C. Kalogeri ⁽¹⁾

Author Affiliations

- 1. Atmospheric Modeling and Weather Forecasting Group, Department of Physics, University of Athens, University Campus, Building Physics V, Athens, Greece

Continue reading...

To view the rest of this content please follow the download PDF link above.

Over 8.5 million scientific documents at your fingertips
© Springer, Part of Springer Science+Business Media

